

A COMPOSITIONAL SHUFFLE CONJECTURE SPECIFYING TOUCH POINTS OF THE DYCK PATH

J. HAGLUND, J. MORSE, AND M. ZABROCKI

ABSTRACT. We introduce a q, t -enumeration of Dyck paths which are forced to touch the main diagonal at specific points and forbidden to touch elsewhere and conjecture that it describes the action of the Macdonald theory ∇ operator applied to a Hall-Littlewood polynomial. Our conjecture refines several earlier conjectures concerning the space of diagonal harmonics including the “shuffle conjecture” (Duke J. Math. **126** (2005), pp. 195 – 232) for $\nabla e_n[X]$. We bring to light that certain generalized Hall-Littlewood polynomials indexed by compositions are the building blocks for the algebraic combinatorial theory of q, t -Catalan sequences and we prove a number of identities involving these functions.

1. INTRODUCTION

Our study concerns the combinatorics behind the character of the space of diagonal harmonics DH_n and identities involving Macdonald polynomials that can be used to form expressions for this character. At the root of this theory is a linear operator ∇ , introduced in [BGHT99], under which the modified Macdonald polynomials $\tilde{H}_\mu[X; q, t]$ are eigenfunctions. [Hai02] proved that the Frobenius image of the character of DH_n equals ∇e_n . This gives an explicit expression involving rational functions in q, t for the multiplicity of an irreducible indexed by a partition λ in the character of DH_n .

An important open problem in this area is the “shuffle conjecture” of [HHLRU05] which asserts that the coefficient of m_λ in ∇e_n simplifies to a q, t statistic on lattice paths. A major breakthrough in this direction was made with the conjectured combinatorial formula of [Hag03] for the coefficient of $m_{1^n} = s_{1^n}$ in ∇e_n . In this case, the coefficient is a bi-graded version of the sign character and it is called the q, t -Catalan $C_n(q, t)$ since it reduces to the n th Catalan number when $q = t = 1$. The combinatorial formula for $C_n(q, t)$ was proven in [GH01], [GH02] and pursuant work [Hag04] also settled the shuffle conjecture for partitions of hook-shape. However, the general case remains a mystery.

An unrelated study of Macdonald polynomials led to the discovery [LLM03] of a new family of symmetric functions called k -Schur functions $s_\lambda^{(k)}[X; t]$ which were conjectured to refine the special combinatorial properties held by Schur functions. The k -Schur functions have a number of conjecturally equivalent characterizations and it has now been established in [LM08, Lam08] that those introduced in [LM07] refine combinatorial, geometric and representation theoretic aspects of Schur theory. This prompted Bergeron, Descouens, and Zabrocki to explore the role of k -Schur

1991 *Mathematics Subject Classification.* Primary 05E05, Secondary 33D52.

Key words and phrases. Dyck Paths, Parking functions, Hall-Littlewood symmetric functions. Work supported by NSF grants DMS-0553619 and DMS-0901467.

functions in the q, t -Catalan theory. To this end, they conjectured in [BDZ10] that the coefficient of s_{1^n} in $\nabla s_{1^n}^{(k)}[X; t]$ is a positive polynomial in q, t and they proved their conjecture for the case that $t = 1$.

Our work here was initially motivated by a desire to find a combinatorial description for this coefficient in general, ideally in terms of a q, t -statistic on lattice paths as with the q, t -Catalans. We found such a description, but more remarkably this led us to discover that a natural setting for the combinatorial theory of DH_n is created by applying ∇ to the general set of Hall-Littlewood polynomials indexed by compositions. To be precise, it was proven in [LM03] that the k -Schur function $s_{1^n}^{(k)}[X; t]$ is merely a certain Hall-Littlewood polynomial. This led us to study ∇ on a Hall-Littlewood polynomial indexed by any partition λ . But in fact, our work carries through to the family of polynomials $C_\alpha[X; q]$, for any composition α , defined in terms of operators similar to Jing operators.

A key component in the proof of the q, t -Catalan conjecture [GH02] is the use of symmetric functions $E_{n,k}[X; q]$ which decompose e_n into pieces that remain positive under the action of ∇ . We have discovered that the $C_\alpha[X; q]$ can be used as building blocks in the q, t -Catalan theory that decompose the $E_{n,k}[X; q]$ into finer pieces, still positive under the action of ∇ . Our conjectures on these building blocks thus refine earlier conjectures involving $E_{n,k}[X; q]$, the conjectures in [BDZ10], the shuffle conjecture, and the conjectures in [BGHT99] asserting that ∇ applied to Hall-Littlewood functions have q, t -positive Schur coefficients. Loehr and Warrington [LW08] introduced an intricate conjecture for the combinatorics of ∇ applied to a Schur function s_λ . Our conjecture is extremely simple, describes the action of ∇ on a larger set of symmetric functions than just a basis, and refines the conjecture of Loehr and Warrington when ∇ acts on the Schur function $s_{(n-k, 1^k)}$ [LW08, Conjecture 3] as explained at the end of Section 4.

Garsia, Xin, and Zabrocki [GXZ10] using work of Hicks [Hic10] have now proven our generalized q, t -Catalan conjecture and expanded the result giving a “compositional q, t -Schröder” theorem.

2. DEFINITIONS AND NOTATION

2.1. Combinatorics. A Dyck path is a lattice path in the first quadrant of the xy -plane from the point $(0, 0)$ to the point (n, n) with steps $+(0, 1)$ and $+(1, 0)$ which stays above the line $x = y$. For a Dyck path D , the cells in the i^{th} row are those unit squares in the xy -plane that are below the path and fully above the line $x = y$ whose NE corner has a y coordinate of i . The set of Dyck paths from $(0, 0)$ to (n, n) will be denoted DP^n and the number of paths in this set is well known to be the Catalan number

$$C_n = \frac{1}{n+1} \binom{2n}{n}.$$

For a Dyck path D , let $a_i = a_i(D)$ equal the number of cells in the i^{th} row of D . It is always true $a_1 = 0$ and $0 \leq a_{i+1} \leq a_i + 1$. We define the arm sequence $\text{arm}(D) = (a_1, a_2, \dots, a_n)$ and note this completely determines D . We consider two statistics (non-negative integers) on Dyck paths. The *area* statistic is the number of whole cells which are below the path and above the diagonal, or


$$\text{area}(D) = \sum_{i=1}^n a_i.$$

The *dinv* statistic is defined as

$$\text{dinv}(D) = \sum_{1 \leq i < j \leq n} \chi(a_i - a_j \in \{0, 1\})$$

where $\chi(\text{true}) = 1$ and $\chi(\text{false}) = 0$.

Remark 2.1. The Dyck path D with arm sequence (0^n) has $\text{area}(D) = 0$ and $\text{dinv}(D) = \binom{n}{2}$. The Dyck path D' with arm sequence $(0, 1, 2, 3, \dots, n-1)$ has $\text{area}(D') = \binom{n}{2}$ and $\text{dinv}(D') = 0$.



Remark 2.2. The original proof of the combinatorial interpretation of the q, t -Catalan was stated in terms of a third statistic $\text{bounce}(D)$. Since we are more cleanly able to formulate our results in terms of the $\text{dinv}(D)$ statistic we choose to state all the results in this paper in terms of the $\text{dinv}(D)$ statistic however the reference [Hag08, p. 50] describes an automorphism ϕ on DP^n such that $\text{area}(\phi(D)) = \text{bounce}(D)$ and $\text{dinv}(\phi(D)) = \text{area}(D)$.

We make use of a partial order on Dyck paths; namely $D_1 \leq D_2$ if $\text{arm}(D_1) \leq \text{arm}(D_2)$, component-wise. In this case we say that D_1 is ‘below’ D_2 because D_1 will not cross D_2 and is hence weakly ‘between’ D_2 and the diagonal.

A composition α of n , denoted $\alpha \models n$, is an integer sequence $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_r)$ with $\alpha_i \geq 1$ and where $|\alpha| = \alpha_1 + \alpha_2 + \dots + \alpha_r = n$. The length of α is $\ell(\alpha) = r$. We shall also use $\overleftarrow{\alpha} = (\alpha_{\ell(\alpha)}, \alpha_{\ell(\alpha)-1}, \dots, \alpha_2, \alpha_1)$. For any composition α , we define

$$(2.1) \quad n(\alpha) = \sum_{i=1}^{\ell(\alpha)} (i-1)\alpha_i.$$

The descent set of a composition α is defined to be

$$\text{Des}(\alpha) = \{\alpha_1, \alpha_1 + \alpha_2, \dots, \alpha_1 + \alpha_2 + \dots + \alpha_{\ell(\alpha)-1}\}.$$

There is a common partial order defined on composition $\alpha, \beta \models n$ by letting $\alpha \leq \beta$ when α is ‘finer’ than β , i.e. $\text{Des}(\beta) \subseteq \text{Des}(\alpha)$. If α is a composition of n , $DP(\alpha)$ represents the Dyck path consisting of α_1 steps in the North $(0, 1)$ direction followed by α_1 steps in the East $(1, 0)$ direction, α_2 $(0, 1)$ steps followed by α_2 $(1, 0)$ steps, etc.

A partition $\lambda = (\lambda_1, \dots, \lambda_r)$ is a non-increasing sequence of positive integers. When λ is a partition of n , denoted $\lambda \vdash n$, $|\lambda| = \sum \lambda_i = n$. The length of λ is $\ell(\lambda) = r$. Given a partition λ , we set

$$(2.2) \quad m(\lambda) = (m_1(\lambda), m_2(\lambda), m_3(\lambda), \dots, m_{|\lambda|}(\lambda)),$$

where the numbers $m_i(\lambda)$ represent the number of parts of size i in λ . The conjugate of a partition λ is the partition $\lambda' = (\lambda'_1, \lambda'_2, \dots, \lambda'_m)$ where λ'_i is the number of parts of λ that are at least i . Partitions are generally considered to be compositions with


parts arranged in non-increasing order. Hence, notions defined on compositions apply to partitions as well. Generally, we will use the symbols α, β, γ to represent compositions and λ, μ, ν for partitions.

For a given Dyck path D , $\text{touch}(D)$ denotes the composition $\gamma = (\gamma_1, \gamma_2, \dots, \gamma_{\ell(\gamma)}) \models n$ that specifies which rows the Dyck path ‘touches’ the diagonal. That is, for $\text{arm}(D) = (a_1, a_2, \dots, a_n)$, $a_k = 0$ if and only if $k = 1$ or $k - 1 \in \text{Des}(\gamma)$. The title of this paper comes from the notion of the touch composition. By requiring that $\text{touch}(D) = \alpha$ for a fixed composition α , we have specified that the Dyck path will touch the diagonal in rows $1, 1 + \alpha_1, 1 + \alpha_1 + \alpha_2, \dots, 1 + \alpha_1 + \alpha_2 + \dots + \alpha_{\ell(\alpha)-1}$ and is forbidden to touch the diagonal in the other rows. Note that under this definition, we view all paths as touching the diagonal in row 1, but none touching in row $n + 1$, and we say the path touches the diagonal $\ell(\alpha)$ times. The partial order on compositions is consistent with the partial order on Dyck paths in the sense that if D_1 and D_2 are Dyck paths such that $D_1 \leq D_2$, then $\text{touch}(D_1) \leq \text{touch}(D_2)$.

Using these notions we introduce a new statistic $\text{doff}_\alpha(D)$ for a given Dyck path D with $\text{touch}(D) \leq \alpha$. If $\text{arm}(D) = (a_1, a_2, \dots, a_n)$, let r_1 be the number of rows such that $a_i = 0$ for $1 \leq i \leq \alpha_1$, r_2 be the number of rows such that $a_i = 0$ for $\alpha_1 < i \leq \alpha_1 + \alpha_2$, and more generally $r_k = \#\{i : a_i = 0 \text{ and } \sum_{j=1}^{k-1} \alpha_j < i \leq \sum_{j=1}^k \alpha_j\}$. We then set

$$(2.3) \quad \text{doff}_\alpha(D) = \sum_{k=1}^{\ell(\alpha)} (\ell(\alpha) - k) r_k .$$

Remark 2.3. If $\text{arm}(D) = (0, 1, 2, 0, 1, 2, 2, 1, 0, 1, 2, 3, 2, 1)$, then $\text{touch}(D) = (3, 5, 6)$. Taking $\alpha = (8, 6)$ we have $\text{doff}_{(8,6)}(D) = 2$.



The only Dyck path with $\text{touch}(D) = (1^n)$ has $\text{arm}(D) = (0^n)$. There are C_{n-1} Dyck paths with $\text{touch}(D) = (n)$ and more generally there are $\prod_{i=1}^{\ell(\alpha)} C_{\alpha_i-1}$ Dyck paths such that $\text{touch}(D) = \alpha$. Note that if D and E are two Dyck paths with $\text{touch}(D) = \text{touch}(E) \leq \alpha$, then $\text{doff}_\alpha(D) = \text{doff}_\alpha(E)$.

The results we mention so far are stated in terms of Dyck paths, but we will require the notion of parking functions to state the generalization of the shuffle conjecture. For a Dyck path D in DP^n with $\text{arm}(D) = (a_1, \dots, a_n)$, let \mathcal{WP}_D be the set of words of length n in the alphabet $\{1, 2, \dots, n\}$ such that $w_i < w_{i+1}$ if $a_i < a_{i+1}$. We use the notation x^w to denote the monomial $x_{w_1} x_{w_2} \dots x_{w_n}$. We

also define an extension of the dinv statistic for words in \mathcal{WP}_D by setting

$$\begin{aligned} \text{dinv}(w) = & |\{(i, j) : 1 \leq i < j \leq n, a_i = a_j \text{ and } w_i < w_j\}| \\ & + |\{(i, j) : 1 \leq i < j \leq n, a_i = a_j + 1 \text{ and } w_i > w_j\}|. \end{aligned}$$

2.2. Symmetric functions. Let X represent a sum of an infinite set of variables $X = x_1 + x_2 + x_3 + \cdots$ considered as elements of the ring of polynomial series in an infinite number of variables of bounded degree. For $r > 0$, let p_r represent a linear and algebraic morphism which acts on polynomial series by $p_r[x] = x^r$. That is for two polynomial series of bounded degree A and B ,

$$\begin{aligned} p_r[A + B] &= p_r[A] + p_r[B] \\ p_r[A - B] &= p_r[A] - p_r[B] \\ p_r[AB] &= p_r[A]p_r[B] \end{aligned}$$

and in particular, $p_r[X] = x_1^r + x_2^r + x_3^r + \cdots$ represents the r^{th} power sum in the variables $\{x_1, x_2, x_3, \dots\}$. The ring of symmetric functions over the field F is defined to be the polynomial ring

$$\Lambda = F[p_1[X], p_2[X], p_3[X], \dots].$$

For our purposes, we choose the field F to be the ring of rational power series in the variables q, t, u, z over \mathbb{Q} where each of the parameters q, t, u, z all have the property that $p_r[a] = a^r$ for each $a = q, t, z, u$.

Generally our symmetric functions f will be considered as polynomials in the elements p_r and then the notation $f[A]$ represents f with each p_r replaced by $p_r[A]$. The degree of p_r is r and the degree of a symmetric function f is determined by the degree of the monomials in the power sums which appear in f . Following the notation of Macdonald [Mac95], we have the power sum basis $p_\lambda[X]$, Schur basis $s_\lambda[X]$, homogeneous basis $h_\lambda[X]$, and elementary basis $e_\lambda[X]$.

In the expressions of variables it is useful to have a special symbol ϵ which will represent a value of negative one but behaves differently than a negative symbol. If f is of homogeneous degree r ,

$$\begin{aligned} f[\epsilon X] &= (-1)^r f[X] \\ f[-\epsilon X] &= \omega(f[X]) \end{aligned}$$

where ω is an involution on symmetric functions such that $\omega(p_\lambda[X]) = (-1)^{|\lambda| + \ell(\lambda)} p_\lambda[X]$, $\omega(e_n[X]) = h_n[X]$ and $\omega(s_\lambda[X]) = s_{\lambda'}[X]$. We will also make use of the standard Hall scalar product which is defined by

$$\langle p_\lambda[X], p_\mu[X]/z_\mu \rangle = \langle s_\lambda[X], s_\mu[X] \rangle = \chi(\lambda = \mu)$$

where $z_\mu = \prod_{i \geq 1} m_i(\mu)! i^{m_i(\mu)}$.

For any symmetric function f , multiplication by f is an operation on symmetric functions which raises the degree of the symmetric function by $\deg(f)$. If we define f^\perp to be the operation which is dual to multiplication in the sense that

$$\langle f^\perp g, h \rangle = \langle g, fh \rangle,$$

then f^\perp is an operator which lowers the degree of the symmetric function by $\deg(f)$. It is not difficult to show that

$$(2.4) \quad f[X + z] = \sum_{k \geq 0} z^k (h_k^\perp f)[X],$$

$$(2.5) \quad f[X - z] = \sum_{k \geq 0} (-z)^k (e_k^\perp f)[X].$$

In addition we will refer to the form of the Macdonald basis $\tilde{H}_\lambda[X; q, t]$ that is relevant to the study of the $n!$ Theorem [Hai01] and the q, t -Catalan numbers. The relation of this basis to the integral form $J_\mu[X; q, t]$ of [Mac95] is

$$\tilde{H}_\mu[X; q, t] = t^{n(\mu)} J_\mu \left[\frac{X}{1 - 1/t}; q, 1/t \right].$$

It is also characterized as the unique basis such that

$$\langle \tilde{H}_\mu[X(1 - 1/t); q, t], \tilde{H}_\lambda[X(1 - q); q, t] \rangle = 0$$

if $\lambda \neq \mu$ and $\langle \tilde{H}_\mu[X; q, t], h_n[X] \rangle = 1$.

We are particularly interested in the Hall-Littlewood symmetric functions. Following the notation of Macdonald we define the functions $Q'_\lambda[X; q]$ to be the basis of the symmetric functions which satisfy

$$\langle Q'_\lambda[X(1 - q); q], Q'_\mu[X; q] \rangle = 0$$

if $\lambda \neq \mu$ and $\langle Q'_\lambda[X; q], h_n[X] \rangle = q^{n(\lambda)}$. Relating the definitions of the Hall-Littlewood and Macdonald symmetric functions, we note that

$$Q'_\lambda[X; q] = \tilde{H}_\lambda[X; 0, 1/q] q^{n(\lambda)} = \sum_{\lambda} K_{\lambda\mu}(q) s_\lambda[X].$$

The operator ∇ was introduced in [BGHT99] and is defined by

$$(2.6) \quad \nabla \tilde{H}_\lambda[X; q, t] = t^{n(\lambda)} q^{n(\lambda')} \tilde{H}_\lambda[X; q, t].$$

This operator has been fundamental to the study of the q, t -combinatorial identities associated with DH_n and Macdonald polynomials. Its definition is chosen so that


$$(2.7) \quad \langle \nabla(e_n[X]), e_n[X] \rangle = C_n(q, t)$$

where $C_n(q, t)$ is the q, t -Catalan polynomial. References [GH01, Hag03, Hag08] showed that

$$(2.8) \quad C_n(q, t) = \sum_{D \in \text{DP}^n} t^{\text{area}(D)} q^{\text{dinv}(D)}$$

with the sum over all Dyck paths of length n .

Remark 2.4. A small example is $C_3(q, t) = q^3 + qt + q^2t + qt^2 + t^3$, whose terms can be computed (in order) from the 5 Dyck paths of length 3 with respective arm sequences $(0, 0, 0)$, $(0, 0, 1)$, $(0, 1, 0)$, $(0, 1, 1)$ and $(0, 1, 2)$:



We will make use of the Newton element

$$(2.9) \quad \Omega[X] = \sum_{\lambda} p_{\lambda}[X]/z_{\lambda} = \sum_{m \geq 0} h_m[X],$$

where we have the identities

$$\begin{aligned} \Omega[X + Y] &= \Omega[X]\Omega[Y] \\ \Omega[X - Y] &= \Omega[X]/\Omega[Y] \\ \Omega[x_1 + \epsilon x_2 - x_3 - \epsilon x_4] &= \frac{(1 - x_3)(1 + x_4)}{(1 - x_1)(1 + x_2)} \\ \Omega[XY] &= \sum_{\lambda} s_{\lambda}[X]s_{\lambda}[Y] = \sum_{\lambda} p_{\lambda}[X]p_{\lambda}[Y]/z_{\lambda}. \end{aligned}$$

Jing [Jin91] introduced a family of operators \mathbb{H}_m indexed by $m \in \mathbb{Z}$ using the following formal power series in the parameter z ,

$$(2.10) \quad \begin{aligned} \mathbb{H}(z)P[X] &= \sum_{m \in \mathbb{Z}} z^m \mathbb{H}_m P[X] := P \left[X - \frac{1 - q}{z} \right] \Omega[zX] \\ &= \sum_{m \in \mathbb{Z}} z^m \sum_{r \geq 0} (-1)^r h_{m+r}[X] e_r[(1 - q)X]^{\perp} P[X]. \end{aligned}$$

He proved that these operators create the Hall-Littlewood polynomials by adding rows.

Proposition 2.5. [Jin91] *For any partition $\lambda = (\lambda_1, \dots, \lambda_{\ell})$,*

$$Q'_{\lambda}[X; q] = \mathbb{H}_{\lambda_1} \mathbb{H}_{\lambda_2} \cdots \mathbb{H}_{\lambda_{\ell}}(1).$$

3. TWO FAMILIES OF HALL-LITTLEWOOD SYMMETRIC FUNCTIONS

Our primary focus is the study of two families of symmetric functions and the combinatorics surrounding them. These functions arise from the following operators \mathbb{B}_m and \mathbb{C}_m , closely related to Jing's \mathbb{H}_m operators from equation (2.10):

$$(3.1) \quad \begin{aligned} \mathbb{B}(z)P[X] &= \sum_{m \in \mathbb{Z}} z^m \mathbb{B}_m P[X] := P \left[X + \epsilon \frac{(1 - q)}{z} \right] \Omega[-\epsilon z X] \\ &= \sum_{m \in \mathbb{Z}} z^m \sum_{r \geq 0} (-1)^r e_{m+r}[X] h_r[X(1 - q)]^{\perp} P[X] \end{aligned}$$

and

$$(3.2) \quad \begin{aligned} \mathbb{C}(z)P[X] &= \sum_{m \in \mathbb{Z}} z^m \mathbb{C}_m P[X] := -q P \left[X + \epsilon \frac{(1 - q)}{z} \right] \Omega[\epsilon(z/q)X] \\ &= \sum_{m \in \mathbb{Z}} (-1/q)^{m-1} z^m \sum_{r \geq 0} q^{-r} h_{m+r}[X] h_r[X(1 - q)]^{\perp} P[X]. \end{aligned}$$

The symmetric functions of particular interest here are those defined, for any composition α , by setting

$$B_{\alpha}[X; q] = \mathbb{B}_{\alpha_{\ell(\alpha)}} \mathbb{B}_{\alpha_{\ell(\alpha)-1}} \cdots \mathbb{B}_{\alpha_1}(1)$$

and

$$C_{\alpha}[X; q] = \mathbb{C}_{\alpha_1} \mathbb{C}_{\alpha_2} \cdots \mathbb{C}_{\alpha_{\ell(\alpha)}}(1).$$

Note that the operators generating B_α and C_α are both indexed by the parts of α , but are applied in reverse order with respect to one another. This is done so that the associated combinatorial and algebraic identities are more uniform.

These operators are related by way of the equation:

$$(3.3) \quad \mathbb{B}(z) = \omega \mathbb{H}(z) \omega$$

and

$$(3.4) \quad \mathbb{C}(z) = (-q) \mathbb{H}^{q \rightarrow 1/q}(-z/q),$$

or equivalently

$$(3.5) \quad \mathbb{C}_m = (-1/q)^{m-1} \mathbb{H}_m^{q \rightarrow 1/q} = (-1/q)^{m-1} \omega \mathbb{B}_m^{q \rightarrow 1/q} \omega.$$

Thus the functions themselves are related as:

$$(3.6) \quad Q'_\lambda[X; q] = \omega B_{\check{\lambda}}[X; q] = (-q)^{\ell(\lambda) - |\lambda|} C_\lambda[X; 1/q].$$

The Jing operators create Hall-Littlewood polynomials indexed by partitions which form a basis for the symmetric function ring. The C_α and B_α symmetric functions are indexed by compositions and are not linearly independent. The equations above detail how Hall-Littlewood symmetric functions are included in these families and therefore, Schur positive expansions of the C_α and B_α hold in certain cases. However, they are not Schur positive in complete generality. The smallest examples that are not uniformly Schur positive or Schur negative are $B_{(3,1)}[X; q]$ and $C_{(1,3)}[X; q]$.

Remark 3.1. Below is a table of the symmetric functions $B_\alpha[X; q]$ and $C_\alpha[X; q]$ for $\alpha \models 4$. Notice that both $B_{(3,1)}[X; q]$ and $C_{(1,3)}[X; q]$ have mixed signs in their coefficients.

$$\begin{aligned} \begin{bmatrix} B_{(1,1,1,1)}[X; q] \\ B_{(1,1,2)}[X; q] \\ B_{(1,2,1)}[X; q] \\ B_{(2,1,1)}[X; q] \\ B_{(1,3)}[X; q] \\ B_{(2,2)}[X; q] \\ B_{(3,1)}[X; q] \\ B_{(4)}[X; q] \end{bmatrix} &= \begin{bmatrix} q^6 & q^3 + q^4 + q^5 & q^2 + q^4 & q + q^2 + q^3 & 1 \\ q^3 & q + q^2 & q & 1 & 0 \\ q^4 & q^2 + q^3 & q^2 & q & 0 \\ q^5 & q^3 + q^4 & q^3 & q^2 & 0 \\ q & 1 & 0 & 0 & 0 \\ q^2 & q & 1 & 0 & 0 \\ q^3 & q^2 & q - 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} s_{(1,1,1,1)}[X; q] \\ s_{(2,1,1)}[X; q] \\ s_{(2,2)}[X; q] \\ s_{(3,1)}[X; q] \\ s_{(4)}[X; q] \end{bmatrix} \\ \\ \begin{bmatrix} C_{(1,1,1,1)}[X; q] \\ C_{(1,1,2)}[X; q] \\ C_{(1,2,1)}[X; q] \\ C_{(2,1,1)}[X; q] \\ C_{(1,3)}[X; q] \\ C_{(2,2)}[X; q] \\ C_{(3,1)}[X; q] \\ C_{(4)}[X; q] \end{bmatrix} &= \begin{bmatrix} 1 & q^{-3} + q^{-2} + q^{-1} & q^{-4} + q^{-2} & q^{-5} + q^{-4} + q^{-3} & q^{-6} \\ 0 & -q^{-3} & -q^{-4} & -q^{-5} - q^{-4} & -q^{-6} \\ 0 & -q^{-2} & -q^{-3} & -q^{-4} - q^{-3} & -q^{-5} \\ 0 & -q^{-1} & -q^{-2} & -q^{-3} - q^{-2} & -q^{-1} \\ 0 & 0 & q^{-3} - q^{-2} & q^{-4} & q^{-5} \\ 0 & 0 & q^{-2} & q^{-3} & q^{-4} \\ 0 & 0 & 0 & q^{-2} & q^{-3} \\ 0 & 0 & 0 & 0 & q^{-3} \end{bmatrix} \begin{bmatrix} s_{(1,1,1,1)}[X; q] \\ s_{(2,1,1)}[X; q] \\ s_{(2,2)}[X; q] \\ s_{(3,1)}[X; q] \\ s_{(4)}[X; q] \end{bmatrix} \end{aligned}$$

To manipulate these symmetric functions, we derive commutation relations between the symmetric function operators. Our first result enables us to expand an element $C_\alpha[X; q]$, for any composition α , in terms of the $C_\lambda[X; q]$ indexed by partitions λ .

Proposition 3.2. *For $m, n \in \mathbb{Z}$, we have*

$$(3.7) \quad q\mathbb{C}_m\mathbb{C}_n - \mathbb{C}_{m+1}\mathbb{C}_{n-1} = \mathbb{C}_n\mathbb{C}_m - q\mathbb{C}_{n-1}\mathbb{C}_{m+1} .$$

In particular for $m \in \mathbb{Z}$,

$$(3.8) \quad \mathbb{C}_m\mathbb{C}_{m+1} = \frac{1}{q}\mathbb{C}_{m+1}\mathbb{C}_m .$$

Proof. We begin with the expressions for the \mathbb{C} -operators from equation (3.2). For ease of notation we shall use h_m in place of $h_m[X]$ and $h_r^{q\perp}$ in place of the expression $h_r[X(1-q)]^\perp$. We compute that

$$(3.9) \quad \begin{aligned} h_r^{q\perp}(h_m P[X]) &= h_m[X + (1-q)z]P[X + (1-q)z] \Big|_{z^r} \\ &= \sum_{i \geq 0} h_{m-i} h_i [1-q] P[X + (1-q)z] \Big|_{z^{r-i}} \\ &= \sum_{i \geq 0} h_{m-i} h_i [1-q] h_{r-i}^{q\perp} P[X]. \end{aligned}$$

We also know that

$$(3.10) \quad h_r[1-q] = \begin{cases} 0 & \text{if } r < 0, \\ 1 & \text{if } r = 0, \\ 1-q & \text{if } r > 0. \end{cases}$$

These two identities imply that,

$$(3.11) \quad \begin{aligned} q\mathbb{C}_m\mathbb{C}_n &= (-1/q)^{m+n-2} \sum_{i \geq 0} \sum_{r \geq 0} \sum_{s \geq 0} q^{-r-s-i+1} h_{m+r+i} h_{n+s-i} h_i [1-q] h_r^{q\perp} h_s^{q\perp} \\ &= (-1/q)^{m+n-2} \sum_{i \geq 0} \sum_{r \geq 0} \sum_{s \geq 0} q^{-r-s-i+1} h_{m+r+i} h_{n+s-i} h_r^{q\perp} h_s^{q\perp} \\ &\quad - (-1/q)^{m+n-2} \sum_{i \geq 1} \sum_{r \geq 0} \sum_{s \geq 0} q^{-r-s-i+2} h_{m+r+i} h_{n+s-i} h_r^{q\perp} h_s^{q\perp} \\ &= (-1/q)^{m+n-2} \sum_{i \geq 0} \sum_{r \geq 0} \sum_{s \geq 0} q^{-r-s-i+1} h_{m+r+i} h_{n+s-i} h_r^{q\perp} h_s^{q\perp} \\ &\quad - (-1/q)^{m+n-2} \sum_{i \geq 0} \sum_{r \geq 0} \sum_{s \geq 0} q^{-r-s-i+1} h_{m+r+i+1} h_{n+s-i-1} h_r^{q\perp} h_s^{q\perp} \\ &= (-1/q)^{m+n-2} \sum_{r \geq 0} \sum_{s \geq 0} \sum_{i \geq 0} q^{-r-s-i+1} s_{(m+r+i, n+s-i)} h_r^{q\perp} h_s^{q\perp}. \end{aligned}$$

where to arrive at (3.11) we have introduced the Schur function $s_{(a,b)} = h_a h_b - h_{a+1} h_{b-1}$. Similarly,

$$\mathbb{C}_{m+1}\mathbb{C}_{n-1} = (-1/q)^{m+n-2} \sum_{r \geq 0} \sum_{s \geq 0} \sum_{i \geq 0} q^{-r-s-i} s_{(m+r+i+1, n+s-i-1)} h_r^{q\perp} h_s^{q\perp} .$$

From these identities, we find that all terms in the difference $q\mathbb{C}_m\mathbb{C}_n - \mathbb{C}_{m+1}\mathbb{C}_{n-1}$ cancel except the $i = 0$ term in (3.11):

$$\begin{aligned} q\mathbb{C}_m\mathbb{C}_n - \mathbb{C}_{m+1}\mathbb{C}_{n-1} &= (-1/q)^{m+n-2} \sum_{r \geq 0} \sum_{s \geq 0} q^{-r-s+1} s_{(m+r, n+s)} h_r^{q\perp} h_s^{q\perp} \\ &\quad + (-1/q)^{m+n-2} \sum_{r \geq 0} \sum_{s \geq 0} \sum_{i \geq 1} q^{-r-s-i+1} s_{(m+r+i, n+s-i)} h_r^{q\perp} h_s^{q\perp} \\ &\quad - (-1/q)^{m+n-2} \sum_{r \geq 0} \sum_{s \geq 0} \sum_{i \geq 0} q^{-r-s-i} s_{(m+r+i+1, n+s-i-1)} h_r^{q\perp} h_s^{q\perp} \\ &= (-1/q)^{m+n-2} \sum_{r \geq 0} \sum_{s \geq 0} q^{-r-s+1} s_{(m+r, n+s)} h_r^{q\perp} h_s^{q\perp}. \end{aligned}$$

We can then compute $\mathbb{C}_n\mathbb{C}_m - q\mathbb{C}_{n-1}\mathbb{C}_{m+1}$ from this by replacing $m \rightarrow n-1$ and $n \rightarrow m+1$. In particular,

$$q\mathbb{C}_{n-1}\mathbb{C}_{m+1} - \mathbb{C}_n\mathbb{C}_m = (-1/q)^{m+n-2} \sum_{r \geq 0} \sum_{s \geq 0} q^{-r-s+1} s_{(n-1+r, m+1+s)} h_r^{q\perp} h_s^{q\perp}.$$

The identity $-s_{(b-1, a+1)} = s_{(a, b)}$ and the commutation of h^\perp then implies our claim. \square

An important consequence of this result is that if α is a composition of length ℓ , then $C_\alpha[X; q]$ can be written as a linear combination of the $C_\lambda[X; q]$ where λ are partitions that also have length ℓ .

Remark 3.3. The symmetric function $C_{(1,3)}[X; q]$ can be expressed in terms of $C_{(3,1)}[X; q]$ and $C_{(2,2)}[X; q]$ using this commutation relation since $q\mathbb{C}_1\mathbb{C}_3 = \mathbb{C}_2\mathbb{C}_2 + \mathbb{C}_3\mathbb{C}_1 - q\mathbb{C}_2\mathbb{C}_2$. Consequently, $C_{(1,3)}[X; q] = (1/q - 1)C_{(2,2)}[X; q] + 1/qC_{(3,1)}[X; q]$.

The relation of \mathbb{C} to \mathbb{B} given in (3.5) enables us to derive an identity on \mathbb{B} from Theorem 3.2. In particular, we simply apply ω to (3.7) and replace q by $1/q$.

Corollary 3.4. For $m \in \mathbb{Z}$,

$$\mathbb{B}_m\mathbb{B}_n - q\mathbb{B}_{m+1}\mathbb{B}_{n-1} = q\mathbb{B}_n\mathbb{B}_m - \mathbb{B}_{n-1}\mathbb{B}_{m+1}.$$

In particular, letting $n = m+1$ gives

$$\mathbb{B}_m\mathbb{B}_{m+1} = q\mathbb{B}_{m+1}\mathbb{B}_m.$$

In fact, we can also pin down commutation relations between the \mathbb{B} and \mathbb{C} operators if $m+n > 0$ (note, the relation does not hold when $m+n \leq 0$).

Proposition 3.5. If $m+n > 0$, then

$$(3.12) \quad \mathbb{B}_n\mathbb{C}_m = q\mathbb{C}_m\mathbb{B}_n.$$

Proof. We use identities (3.9) and (3.10) and compute an expression for $\mathbb{B}_m\mathbb{C}_n$:

$$\begin{aligned} \mathbb{B}_m\mathbb{C}_n &= (-1/q)^{n-1} \sum_{r \geq 0} \sum_{s \geq 0} \sum_{i \geq 0} (-1)^r q^{-s} e_{m+r} h_{n+s-i} h_i [1-q] h_{r-i}^{q\perp} h_s^{q\perp} \\ &= (-1/q)^{n-1} \sum_{r \geq 0} \sum_{s \geq 0} (-1)^r q^{-s} e_{m+r} h_{n+s} h_r^{q\perp} h_s^{q\perp} \\ &\quad + (-1/q)^{n-1} \sum_{i \geq 1} \sum_{r \geq 0} \sum_{s \geq 0} (-1)^{r+i} q^{-s} e_{m+r+i} h_{n+s-i} (1-q) h_r^{q\perp} h_s^{q\perp}. \end{aligned}$$

Analogously, we also have the equations $h_r^{q\perp} e_m = \sum_{i \geq 0} e_{m-i} e_i [1 - q] h_{r-i}^{q\perp}$ and $e_r [1 - q] = (-q)^{r-1} h_r [1 - q]$ if $r > 0$. From this we derive a similar expression for $q\mathbb{C}_n \mathbb{B}_m$:

$$\begin{aligned} q\mathbb{C}_n \mathbb{B}_m &= (-1/q)^{n-1} \sum_{s \geq 0} \sum_{r \geq 0} \sum_{i \geq 0} (-1)^r q^{-s+1} h_{n+s} e_{m+r-i} e_i [1 - q] h_{s-i}^{q\perp} h_r^{q\perp} \\ &= (-1/q)^{n-1} \sum_{s \geq 0} \sum_{r \geq 0} (-1)^r q^{-s+1} e_{m+r} h_{n+s} h_s^{q\perp} h_r^{q\perp} \\ &\quad + (-1/q)^{n-1} \sum_{i \geq 1} \sum_{s \geq 0} \sum_{r \geq 0} (-1)^{r+i+1} q^{-s} e_{m+r-i} h_{n+s+i} (1 - q) h_s^{q\perp} h_r^{q\perp}. \end{aligned}$$

Their difference is

$$\begin{aligned} \mathbb{B}_m \mathbb{C}_n - q\mathbb{C}_n \mathbb{B}_m &= (1 - q)(-1/q)^{n-1} \sum_{s \geq 0} \sum_{r \geq 0} q^{-s} (-1)^r e_{m+r} h_{n+s} h_r^{q\perp} h_s^{q\perp} \\ &\quad + (1 - q)(-1/q)^{n-1} \sum_{r \geq 0} \sum_{s \geq 0} q^{-s} \left(\sum_{i \geq 1} (-1)^{r+i} e_{m+r-i} h_{n+s+i} \right) h_r^{q\perp} h_s^{q\perp} \\ &\quad + (1 - q)(-1/q)^{n-1} \sum_{r \geq 0} \sum_{s \geq 0} q^{-s} \left(\sum_{i \geq 1} (-1)^{r+i} e_{m+r+i} h_{n+s-i} \right) h_r^{q\perp} h_s^{q\perp}. \end{aligned}$$

In fact, the right hand side reduces to zero since $m + n > 0$ implies that for each $r, s \geq 0$, $m + n + r + s > 0$ and

$$\begin{aligned} &(-1)^r e_{m+r} h_{n+s} + \sum_{i \geq 1} (-1)^{r+i} e_{m+r-i} h_{n+s+i} + \sum_{i \geq 1} (-1)^{r+i} e_{m+r+i} h_{n+s-i} \\ &= \sum_{i=-n-s}^{m+r} (-1)^{r+i} e_{m+r-i} h_{n+s+i} = 0 \end{aligned}$$

by the identity $\sum_{i=0}^d (-1)^i e_{d-i} h_i = 0$ for all $d > 0$. \square

Jing's operators generalize operators of Bernstein (see [Mac95]), defined by

$$\begin{aligned} (3.13) \quad \mathbb{S}(z)P[X] &= \sum_{m \in \mathbb{Z}} z^m \mathbb{S}_m P[X] = P \left[X - \frac{1}{z} \right] \Omega[zX] \\ &= \sum_{m \in \mathbb{Z}} z^m \sum_{r \geq 0} (-1)^r h_{m+r}[X] e_r[X]^\perp P[X]. \end{aligned}$$

These are creation operators for the Schur functions since $\mathbb{S}_{\lambda_1} \mathbb{S}_{\lambda_2} \cdots \mathbb{S}_{\lambda_\ell}(1) = s_\lambda[X]$, and they satisfy the commutation relation $\mathbb{S}_m \mathbb{S}_n = -\mathbb{S}_{n-1} \mathbb{S}_{m+1}$. We can write the Schur creation operators in terms of the \mathbb{C}_a operators, which will help us in § 5 to write Schur functions in terms of the \mathbb{C}_α .

Proposition 3.6. *For $m \in \mathbb{Z}$,*

$$(3.14) \quad \mathbb{S}_m = (-q)^{m-1} \sum_{i \geq 0} \mathbb{C}_{m+i} e_i^\perp.$$

Proof. We will use the identity $h_r[(1-q)X] = \sum_{j \geq 0} h_j[X] h_{r-j}[-qX] = \sum_{j \geq 0} (-q)^{r-j} h_j[X] e_{r-j}[X]$ and calculate directly,

$$\begin{aligned}
(-q)^{m-1} \sum_{i \geq 0} \mathbb{C}_{m+i} e_i^\perp &= (-q)^{m-1} \sum_{i \geq 0} (-1/q)^{m+i-1} \sum_{r \geq 0} q^{-r} h_{m+i+r}[X] h_r[X(1-q)]^\perp e_i^\perp \\
&= \sum_{r \geq 0} \sum_{i \geq 0} \sum_{j \geq 0} (-1/q)^{i-r+j} q^{-r} h_{m+i+r}[X] h_j^\perp e_{r-j}^\perp e_i^\perp \\
&= \sum_{d \geq 0} \sum_{j=0}^d \sum_{r \geq 0} (-1)^{d-r} q^{-d} h_{m+d+r-j}[X] h_j^\perp e_{r-j}^\perp e_{d-j}^\perp \\
&= \sum_{d \geq 0} \sum_{j=0}^d \sum_{r \geq 0} (-1)^{d-r-j} q^{-d} h_{m+d+r}[X] h_j^\perp e_r^\perp e_{d-j}^\perp \\
&= \sum_{r \geq 0} (-1)^{-r} h_{m+r}[X] e_r^\perp = \mathbb{S}_m
\end{aligned}$$

where the last equality follows because $\sum_{j=0}^d (-1)^j h_j e_{d-j} = 0$ for all $d > 0$ so the remaining sum is only the part where $d = 0$. \square

Remark 3.7. In the reference [GXZ10], the operator \mathbb{C}_a is presented in a slightly different but equivalent expression. We note that a series $f(z) = \sum_{n \in \mathbb{Z}} f_n z^n$ has the property that $f(\epsilon z/q) \Big|_{z^a} = (-1/q)^a f(z) \Big|_{z^a}$. For this reason,

$$\begin{aligned}
\mathbb{C}_a P[X] &= -qP \left[X + \epsilon \frac{(1-q)}{z} \right] \Omega[\epsilon(z/q)X] \Big|_{z^a} \\
&= -q(-1/q)^a P \left[X + \frac{(1-q)}{qz} \right] \Omega[zX] \Big|_{z^a} \\
&= (-1/q)^{a-1} P \left[X - \frac{1-1/q}{z} \right] \Omega[zX] \Big|_{z^a}.
\end{aligned}$$

4. THE COMBINATORICS OF ∇ APPLIED TO HALL-LITTLEWOOD POLYNOMIALS

Recall that in the special case that α is a partition, $B_\alpha[X; q]$ and $C_\alpha[X; q]$ are closely related to the Hall-Littlewood symmetric functions. It was conjectured in [BGHT99, Conjecture II and III] (partially attributed to A. Lascoux) that applying ∇ to a Hall-Littlewood polynomial produces a Schur positive function. Our main discovery is that including all compositions α in the study of $\nabla(B_\alpha[X; q])$ and $\nabla(C_\alpha[X; q])$ leads to a natural refinement for the combinatorics of Dyck paths. Moreover, our combinatorial exploration led us to discover new symmetric function identities.

One useful tool in the exploration of the operator ∇ is the fact that $\nabla^{q=1}$ is a multiplicative operator. Since we can deduce from the operator definitions of our symmetric functions that $B_\alpha[X; 1] = e_\alpha[X]$ and $C_\alpha[X; 1] = h_\alpha[X]$, we have

$$\nabla^{q=1}(B_\alpha[X; 1]) = \nabla^{q=1}(e_{\alpha_\ell}[X]) \nabla^{q=1}(e_{\alpha_{\ell-1}}[X]) \cdots \nabla^{q=1}(e_{\alpha_1}[X])$$

and

$$\nabla^{q=1}(C_\alpha[X; 1]) = \nabla^{q=1}(h_{\alpha_1}[X]) \nabla^{q=1}(h_{\alpha_2}[X]) \cdots \nabla^{q=1}(h_{\alpha_\ell}[X]).$$

From this we can deduce the coefficient of $e_n[X]$. In particular, the coefficient of $e_n[X]$ in $\nabla(e_n[X])$ and in $\nabla(h_n[X])$ is the q, t -Catalan numbers $C_n(q, t)$ and

$C_{n-1}(q, t)$, respectively. Thus, the coefficient of $e_n[X]$ in $\nabla^{q=1}(B_\alpha[X; 1])$ and in $\nabla^{q=1}(C_\alpha[X; 1])$ is $\prod_i C_{\alpha_i}(1, t)$ and $\prod_i C_{\alpha_i-1}(1, t)$ respectively. The combinatorial interpretation for $C_n(1, t)$ then gives combinatorial meaning to these coefficients. Namely, $\langle e_n[X], \nabla^{q=1}(B_\alpha[X; 1]) \rangle$ is the t -enumeration of Dyck paths (with weight t raised to the area) which lie below the staircase consisting of α_1 steps up and over, α_2 steps up and over, etc. and $\langle e_n[X], \nabla^{q=1}(C_\alpha[X; 1]) \rangle$ is a t -enumeration of Dyck paths which touch the diagonal only in rows $1, 1 + \alpha_1, 1 + \alpha_1 + \alpha_2$ steps, etc.

Proposition 4.1. *For α a composition of n ,*

$$(4.1) \quad \langle \nabla^{q=1}(B_\alpha[X; 1]), e_n[X] \rangle = \sum_{D \leq DP(\alpha)} t^{\text{area}(D)}$$

and

$$(4.2) \quad \langle \nabla^{q=1}(C_\alpha[X; 1]), e_n[X] \rangle = \sum_{\text{touch}(D)=\alpha} t^{\text{area}(D)}.$$

Remarkably, we have empirical evidence to suggest that in general, there is a combinatorial interpretation for the coefficient of $e_n[X]$ in $\nabla(B_\alpha[X; q])$ and in $\nabla(C_\alpha[X; q])$ that naturally generalizes the beautiful combinatorics of the q, t -Catalan.

Conjecture 4.2. *For $\alpha \models n$,*

$$(4.3) \quad \langle \nabla(B_\alpha[X; q]), e_n[X] \rangle = \sum_{D \leq DP(\alpha)} t^{\text{area}(D)} q^{\text{dinv}(D) + \text{doff}_\alpha(D)}.$$

Conjecture 4.3. *For $\alpha \models n$,*

$$(4.4) \quad \langle \nabla(C_\alpha[X; q]), e_n[X] \rangle = \sum_{\text{touch}(D)=\alpha} t^{\text{area}(D)} q^{\text{dinv}(D)}.$$

Our work was inspired by the work of [BDZ10] where they considered coefficients $\nabla(B_{\vec{\lambda}}[X; q])$ for $\vec{\lambda}$ a hook partition (since for that case $\text{doff}_{\vec{\lambda}}(D) = 0$). The innovation in these identities is to consider symmetric functions indexed by compositions which allowed us to conjecture the action of ∇ on a spanning set of the symmetric functions.

More generally, we have conjectures for the expansion of $\nabla(B_\alpha[X; q])$ and $\nabla(C_\alpha[X; q])$ into monomials.

Conjecture 4.4.

$$(4.5) \quad \nabla(B_\alpha[X; q]) = \sum_{D \leq DP(\alpha)} \sum_{w \in WP_D} t^{\text{area}(w)} q^{\text{dinv}(w) + \text{doff}_\alpha(D)} x^w.$$

Conjecture 4.5.

$$(4.6) \quad \nabla(C_\alpha[X; q]) = \sum_{\text{touch}(D)=\alpha} \sum_{w \in WP_D} t^{\text{area}(D)} q^{\text{dinv}(w)} x^w.$$

By the arguments in [HHLRU05] (see also [Hag08, p.99]) Conjecture 4.4 and 4.5 imply Conjecture 4.2 and 4.3. The case $\alpha = (n)$ of (4.5) reduces to the shuffle conjecture since $B_{(n)}[X; q] = e_n[X]$. Also, because of the expansion of $s_{(n-k, 1^k)}$ in Proposition 5.3 in the next section, (4.6) implies the special case of the Loehr-Warrington conjecture [LW08, Conjecture 3] involving the action of ∇ on the Schur function $s_{(n-k, 1^k)}$.

We will prove in the next section that Conjecture 4.4 and 4.5 are equivalent to each other (and by consequence Conjecture 4.2 and 4.3 are equivalent as well). In work building on our results here, [GXZ10] with contributions from [Hic10] proved Conjecture 4.3.

5. SYMMETRIC FUNCTION IDENTITIES

The exploration of q, t -Catalans led [GH01] to the special symmetric function elements $E_{n,k}[X; q]$, defined by the algebraic identity

$$e_n \left[X \frac{1-z}{1-q} \right] = \sum_{k=1}^n \frac{(z; q)_k}{(q; q)_k} E_{n,k}[X; q]$$

where $(z; q)_k = (1-z)(1-qz) \cdots (1-q^{k-1}z)$. These elements play a fundamental role in the proof that the q, t -Catalan polynomial is the q, t -enumeration of Dyck paths as given in (2.8). Namely, the proof follows by showing that

$$\langle \nabla E_{n,k}[X; q], e_n[X] \rangle = q^{\binom{k}{2}} t^{n-k} \sum_{r=0}^{n-k} \left[\begin{matrix} r+k-1 \\ r \end{matrix} \right]_q \langle \nabla(E_{n-k,r}[X; q]), e_{n-r}[X] \rangle ,$$

where $\left[\begin{matrix} n \\ k \end{matrix} \right]_q = \frac{(q; q)_n}{(q; q)_k (q; q)_{n-k}}$, and the combinatorial interpretation for $\langle \nabla E_{n,k}[X; q], e_n[X] \rangle$ in terms of Dyck paths satisfies the same recurrence.

In particular, the coefficient of $e_n[X]$ in $\nabla(E_{n,k}[X; q])$ q, t -enumerates the Dyck paths which touch the diagonal k times. From this, Conjecture 4.3 leads us to expect that

$$\langle e_n[X], \nabla(E_{n,k}[X; q]) \rangle = \sum_{\alpha \models n, \ell(\alpha)=k} \langle e_n[X], \nabla(C_\alpha[X; q]) \rangle .$$

In fact, we have discovered much more generally that

$$(5.2) \quad E_{n,k}[X; q] = \sum_{\substack{\alpha \models n \\ \ell(\alpha)=k}} C_\alpha[X; q] .$$

This section is devoted to proving this surprising result, which suggests that the $C_\alpha[X; q]$ are the building blocks in q, t -Catalan theory.

Remark 5.1. A key step in the proof of our Conjectures 4.2 and 4.3 relies on extending the recurrence (5.1) to involve Dyck paths which touch the diagonal at certain points. This is carried out in [GXZ10].

Our point of departure is to give a simple expression for $e_n[X]$ in terms of the Hall-Littlewood symmetric functions $C_\alpha[X; q]$.

Proposition 5.2.

$$(5.3) \quad e_n[X] = \sum_{\alpha \models n} C_\alpha[X; q] .$$

Proof. Assume by induction on n that equation (5.3) holds (the base cases of $n = 0$ and 1 are easily verified). Since the operator \mathbb{S}_m is a creation operator for the Schur

functions, by (3.14) we have

$$e_n[X] = s_{(1^n)}[X] = \mathbb{S}_1(s_{(1^{n-1})}[X]) = \sum_{i=0}^{n-1} \mathbb{C}_{1+i} s_{(1^{n-i-1})}[X]$$

Which, by induction, gives

$$e_n[X] = \sum_{i=0}^{n-1} \sum_{\alpha \models n-i-1} \mathbb{C}_{i+1} C_\alpha[X; q] = \sum_{\alpha \models n} C_\alpha[X; q] .$$

□

Proposition 5.2 can be stated in a more general form, suggesting that any Schur function may expand nicely in terms of our Hall-Littlewood spanning set.

Proposition 5.3. *For $0 \leq k < n$,*

$$s_{(n-k, 1^k)}[X] = (-q)^{n-k-1} \sum_{\substack{\alpha \models n \\ \alpha_1 \geq n-k}} C_\alpha[X; q] .$$

Proof. Again using that \mathbb{S} is a Schur function creation operator, by (3.14) we have

$$s_{(n-k, 1^k)}[X] = \mathbb{S}_{n-k}(s_{1^k}[X]) = (-q)^{n-k-1} \sum_{i=0}^k \mathbb{C}_{n-k+i} (s_{(1^{k-i})}[X]) .$$

The previous proposition then implies

$$\begin{aligned} s_{(n-k, 1^k)}[X] &= (-q)^{n-k-1} \sum_{i=0}^k \mathbb{C}_{n-k+i} \left(\sum_{\alpha \models k-i} C_\alpha[X; q] \right) \\ &= (-q)^{n-k-1} \sum_{i=0}^k \sum_{\alpha \models k-i} C_{(n-k+i, \alpha)}[X; q] = (-q)^{n-k-1} \sum_{\substack{\alpha \models n \\ \alpha_1 \geq n-k}} C_\alpha[X; q] . \end{aligned}$$

□

We are now in the position to prove that $E_{n,k}[X; q]$ can be decomposed canonically in terms of the $C_\alpha[X; q]$.

Proposition 5.4. *For $0 \leq k < n$,*

$$(5.4) \quad E_{n,k}[X; q] = \sum_{\substack{\mu \vdash n \\ \ell(\mu) = k}} q^{-n(\mu) - k + M(\mu)} \left[\begin{matrix} k \\ m(\mu) \end{matrix} \right]_q C_\mu[X; q],$$

where $M(\mu) = \sum_{i=1}^n \binom{m_i(\mu)+1}{2}$ and $\left[\begin{matrix} k \\ m(\mu) \end{matrix} \right]_q = (q; q)_k / \prod_{i=1}^n (q; q)_{m_i(\mu)}$.

Proof. Recall the expansion of the elementary symmetric functions in the Macdonald basis is given by (see [GH01])

$$e_n \left[X \frac{1-z}{1-q} \right] = \sum_{\mu \vdash n} \frac{\tilde{H}_\mu[X; q, t] \tilde{H}_\mu[(1-z)(1-t); q, t]}{\tilde{h}_\mu(q, t) \tilde{h}'_\mu(q, t)},$$

where $\tilde{h}_\mu(q, t) = \prod_{c \in \mu} (q^{a(c)} - t^{l(c)+1})$ and $\tilde{h}'_\mu(q, t) = \prod_{c \in \mu} (t^{l(c)} - q^{a(c)+1})$. When $t = 0$, these expressions reduce to

$$\tilde{h}_\mu(q, 0) = \prod_{c \in \mu} (q^{a(c)} - 0^{l(c)+1}) = q^{n(\mu')}$$

and

$$\begin{aligned} \tilde{h}'_\mu(q, 0) &= \prod_{\substack{c \in \mu \\ l(c)=0}} (1 - q^{a(c)+1}) \prod_{\substack{c \in \mu \\ l(c) \neq 0}} (-q^{a(c)+1}) \\ &= (-1)^{n-\mu_1} q^{n+n(\mu')-M(\mu')} \prod_{i=1}^n (q; q)_{m_i(\mu')}. \end{aligned}$$

Therefore since $\tilde{H}_\mu[X; q, t] = \tilde{H}_{\mu'}[X; t, q]$, when we set $t = 0$ everywhere we have

$$(5.5) \quad e_n \left[X \frac{1-z}{1-q} \right] = \sum_{\mu \vdash n} (-1)^{n-\mu_1} q^{-n-2n(\mu')+M(\mu')} \frac{\tilde{H}_{\mu'}[X; 0, q] \tilde{H}_\mu[(1-z); q, 0]}{\prod_{i=1}^n (q; q)_{m_i(\mu')}}.$$

Now the evaluation

$$(5.6) \quad \tilde{H}_\mu[(1-z); q, t] = \prod_{c \in \mu} (1 - zt^{l'(c)} q^{a'(c)})$$

also yields

$$\tilde{H}_\mu[(1-z); q, 0] = (z; q)_{\mu_1}.$$

Thus, replacing μ by μ' in (5.5), and thereby exchanging μ_1 and $\ell(\mu)$, gives

$$(5.7) \quad e_n \left[X \frac{1-z}{1-q} \right] = \sum_{\mu \vdash n} (-1)^{n-\ell(\mu)} q^{-n-2n(\mu)+M(\mu)} \frac{\tilde{H}_\mu[X; 0, q] (z; q)_{\ell(\mu)}}{\prod_{i=1}^n (q; q)_{m_i(\mu)}}.$$

Since $\tilde{H}_\mu[X; 0, q] = (-1)^{n-\ell(\mu)} q^{n(\mu)+n-\ell(\mu)} C_\mu[X; q]$, we have

$$e_n \left[X \frac{1-z}{1-q} \right] = \sum_{k=1}^n \frac{(z; q)_k}{(q; q)_k} \sum_{\substack{\mu \vdash n \\ \ell(\mu)=k}} q^{-k-n(\mu)+M(\mu)} \frac{C_\mu[X; q] (q; q)_k}{\prod_{i=1}^n (q; q)_{m_i(\mu)}},$$

which implies our claim. \square

The q -binomial coefficients that appear in equation (5.4) suggest that there is a relation between the terms of the $C_\lambda[X; q]$ basis and subsets of a k element set. It turns out that Proposition 5.4 can be more cleanly written over compositions using a different expansion.

Corollary 5.5. *For $0 \leq k < n$,*

$$(5.8) \quad E_{n,k}[X; q] = \sum_{\substack{\alpha \models n \\ \ell(\alpha)=k}} C_\alpha[X; q].$$

Proof. Using the straightening relations of the \mathbb{C}_m operators, if α is a composition of n and λ is a partition of n such that $\ell(\lambda) \neq \ell(\alpha)$, then

$$C_\alpha[X; q] \Big|_{C_\lambda[X; q]} = 0.$$

Now for $\ell(\lambda) = k$, by Proposition 5.4 and the fact that $e_n[X] = \sum_{k=1}^n E_{n,k}[X; q]$,

$$\begin{aligned} E_{n,k}[X; q] \Big|_{C_\lambda[X; q]} &= e_n[X] \Big|_{C_\lambda[X; q]} \\ &= \sum_{\alpha \models n} C_\alpha[X; q] \Big|_{C_\lambda[X; q]} \\ &= \sum_{\substack{\alpha \models n \\ \ell(\alpha) = k}} C_\alpha[X; q] \Big|_{C_\lambda[X; q]}. \end{aligned}$$

Furthermore if $\ell(\lambda) \neq k$ then

$$E_{n,k}[X; q] \Big|_{C_\lambda[X; q]} = 0 = \sum_{\substack{\alpha \models n \\ \ell(\alpha) = k}} C_\alpha[X; q] \Big|_{C_\lambda[X; q]}.$$

Since the functions $C_\lambda[X; q]$ are a basis this implies that $E_{n,k}[X; q] = \sum_{\substack{\alpha \models n \\ \ell(\alpha) = k}} C_\alpha[X; q]$. \square

We have seen in (3.5) that \mathbb{C} is naturally related to \mathbb{B} . Here we pin down the relationship between the symmetric functions $B_\alpha[X; q]$ and $C_\alpha[X; q]$. A by-product of this identity is that Conjecture 4.5 implies Conjecture 4.4.

Theorem 5.6. *For $n \geq 0$ and any composition $\alpha \models n$,*

$$(5.9) \quad B_\alpha[X; q] = \sum_{\beta \leq \alpha} q^{\text{doff}_\alpha(DP(\beta))} C_\beta[X; q].$$

Proof. We show this result by induction on the number of parts of α . The base case follows since $\mathbb{B}_m(1) = e_m[X]$ which is equal to $\sum_{\gamma \models m} C_\gamma[X; q]$ by Proposition 5.2. Assume by induction that (5.9) holds for a composition α of length ℓ and consider a composition (m, α) . We then have

$$(5.10) \quad B_{(\alpha, m)}[X; q] = \mathbb{B}_m(B_\alpha[X; q]) = \sum_{\beta \leq \alpha} q^{\text{doff}_\alpha(DP(\beta))} \mathbb{B}_m(C_\beta[X; q]).$$

Now consider $\mathbb{B}_m(C_\beta[X; q]) = \mathbb{B}_m \circ \mathbb{C}_{\beta_1} \circ \mathbb{C}_{\beta_2} \circ \dots \circ \mathbb{C}_{\beta_{\ell(\beta)}}(1)$. The commutation relation between the \mathbb{C}_n and \mathbb{B}_m from Theorem 3.5 implies

$$\mathbb{B}_m(C_\beta[X; q]) = q^{\ell(\beta)} \mathbb{C}_{\beta_1} \circ \mathbb{C}_{\beta_2} \circ \dots \circ \mathbb{C}_{\beta_{\ell(\beta)}} \circ \mathbb{B}_m(1).$$

By Proposition 5.2, we then have

$$(5.11) \quad \mathbb{B}_m(C_\beta[X; q]) = q^{\ell(\beta)} \mathbb{C}_{\beta_1} \circ \mathbb{C}_{\beta_2} \circ \dots \circ \mathbb{C}_{\beta_{\ell(\beta)}} \left(\sum_{\gamma \models m} C_\gamma[X; q] \right)$$

$$(5.12) \quad = q^{\ell(\beta)} \sum_{\gamma \models m} C_{(\beta, \gamma)}[X; q].$$

Putting (5.12) into (5.10), we thus find that

$$B_{(\alpha, m)}[X; q] = \sum_{\beta \leq \alpha} q^{\text{doff}_\alpha(DP(\beta)) + \ell(\beta)} \sum_{\gamma \models m} C_{(\beta, \gamma)}[X; q].$$

For each term in the sum, the composition (β, γ) is finer than the composition (α, m) . Moreover, if we let r_i be the number of times that $DP(\beta)$ touches the

diagonal below the i^{th} bump of the Dyck path $DP(\alpha)$, then $\text{doff}_{(\alpha, m)}(DP(\beta, \gamma)) = \sum_{i=1}^{\ell(\alpha)} r_i(\ell(\alpha) + 1 - i) = \sum_{i=1}^{\ell(\alpha)} r_i(\ell(\alpha) - i) + \sum_{i=1}^{\ell(\alpha)} r_i = \text{doff}_{\alpha}(DP) + \ell(\beta)$. \square

To prove that Conjectures 4.5 and 4.4 are in fact equivalent, we need to express $C_{\alpha}[X; q]$ in terms of $B_{\beta}[X; q]$.

Lemma 5.7. *Let γ, α be compositions with $\gamma \leq \alpha$. Then*

$$\sum_{\substack{\beta \\ \gamma \leq \beta \leq \alpha}} (-1)^{\ell(\alpha) - \ell(\beta)} q^{\ell(\alpha) - \ell(\beta) + \text{doff}_{\beta}(DP(\gamma)) - \text{doff}_{\overleftarrow{\alpha}}(DP(\overleftarrow{\beta}))} = \begin{cases} 0 & \text{if } \gamma < \alpha \\ 1 & \text{if } \gamma = \alpha \end{cases}.$$

Proof. First assume that $\gamma < \alpha$ and consider the difference of the descent sets $\text{Des}(\gamma) \setminus \text{Des}(\alpha) = \{i_1, i_2, \dots, i_d\}$. There are 2^d compositions β such that $\gamma \leq \beta \leq \alpha$ and we will pair them up with a sign reversing involution.

If $i_1 \in \text{Des}(\beta)$, let $\tilde{\beta}$ be the composition with $\text{Des}(\tilde{\beta}) = \text{Des}(\beta) \setminus \{i_1\}$ (the terms with $i_1 \in \text{Des}(\beta)$ will match with the terms $i_1 \notin \text{Des}(\tilde{\beta})$). There is some $r > 1$ such that $\alpha_1 = \beta_1$, $\alpha_2 = \beta_2, \dots, \alpha_r = \beta_r$ because $\text{Des}(\beta)$ contains the descent i_1 that is not in $\text{Des}(\alpha)$. Calculating directly we have that,

$$\begin{aligned} \ell(\alpha) - \ell(\beta) &= \ell(\alpha) - \ell(\tilde{\beta}) - 1 \\ \text{doff}_{\beta}(DP(\gamma)) &= \text{doff}_{\tilde{\beta}}(DP(\gamma)) + r \\ -\text{doff}_{\overleftarrow{\alpha}}(DP(\overleftarrow{\beta})) &= -\text{doff}_{\overleftarrow{\alpha}}(DP(\overleftarrow{\tilde{\beta}})) - r + 1. \end{aligned}$$

Therefore

$$\ell(\alpha) - \ell(\beta) + \text{doff}_{\beta}(DP(\gamma)) - \text{doff}_{\overleftarrow{\alpha}}(DP(\overleftarrow{\beta})) = \ell(\alpha) - \ell(\tilde{\beta}) + \text{doff}_{\tilde{\beta}}(DP(\gamma)) - \text{doff}_{\overleftarrow{\alpha}}(DP(\overleftarrow{\tilde{\beta}}))$$

and the signs of these terms are different in the sum. This provides a sign reversing involution and hence those terms with $\text{touch}(D) < \alpha$ sum to 0 matching those terms with $\text{Des}(\beta)$ which include the smallest descent.

Now for those terms with $\text{touch}(D) = \alpha$ we have $\gamma = \beta = \alpha$ and

$$\ell(\alpha) - \ell(\beta) + \text{doff}_{\beta}(DP(\gamma)) - \text{doff}_{\overleftarrow{\alpha}}(DP(\overleftarrow{\beta})) = 0$$

$$\text{since } \text{doff}_{\alpha}(DP(\alpha)) = \binom{\ell(\alpha)}{2}.$$

\square

Theorem 5.8. *For $n \geq 0$ and for any composition $\alpha \models n$,*

$$(5.13) \quad C_{\alpha}[X; q] = \sum_{\beta \leq \alpha} (-q)^{\ell(\alpha) - \ell(\beta)} q^{-\text{doff}_{\overleftarrow{\alpha}}(DP(\overleftarrow{\beta}))} B_{\beta}[X; q].$$

Proof. By Theorem 5.6 we have

$$\begin{aligned} & \sum_{\beta \leq \alpha} (-q)^{\ell(\alpha) - \ell(\beta)} q^{-\text{doff}_{\overleftarrow{\alpha}}(DP(\overleftarrow{\beta}))} B_{\beta}[X; q] \\ &= \sum_{\beta \leq \alpha} (-q)^{\ell(\alpha) - \ell(\beta)} q^{-\text{doff}_{\overleftarrow{\alpha}}(DP(\overleftarrow{\beta}))} \sum_{\gamma \leq \beta} q^{\text{doff}_{\beta}(DP(\gamma))} C_{\gamma}[X; q] \\ &= \sum_{\gamma} C_{\gamma}[X; q] \sum_{\substack{\beta \\ \gamma \leq \beta \leq \alpha}} (-q)^{\ell(\alpha) - \ell(\beta)} q^{-\text{doff}_{\overleftarrow{\alpha}}(DP(\overleftarrow{\beta}))} q^{\text{doff}_{\beta}(DP(\gamma))}, \end{aligned}$$

and our claim now follows by Lemma 5.7. \square

Theorem 5.9. *Conjecture 4.5 is true if and only if Conjecture 4.4 is true.*

Proof. Theorem 5.6 gives $B_\alpha[X; q]$ in terms of $C_\beta[X; q]$, to which we apply ∇ :

$$\nabla(B_\alpha[X; q]) = \sum_{\beta \leq \alpha} q^{\text{doff}_\alpha(DP(\beta))} \nabla(C_\beta[X; q]).$$

Given that Conjecture 4.5 holds, we then have that

$$\begin{aligned} \nabla(B_\alpha[X; q]) &= \sum_{\beta \leq \alpha} \sum_{\substack{D \\ \text{touch}(D)=\beta}} \sum_{w \in WP_D} t^{\text{area}(D)} q^{\text{dinv}(w) + \text{doff}_\alpha(DP(\beta))} x^w \\ &= \sum_{D \leq DP(\beta)} \sum_{w \in WP_D} t^{\text{area}(D)} q^{\text{dinv}(w) + \text{doff}_\alpha(DP(\beta))} x^w. \end{aligned}$$

On the other hand, assuming Conjecture 4.4 holds, Theorem 5.8 gives

$$\begin{aligned} \nabla(C_\alpha[X; q]) &= \sum_{\beta \leq \alpha} (-q)^{\ell(\alpha) - \ell(\beta)} q^{-\text{doff}_{\overleftarrow{\alpha}}(DP(\overleftarrow{\beta}))} \nabla(B_\beta[X; q]) \\ (5.14) \quad &= \sum_{\beta \leq \alpha} \sum_{D \leq DP(\beta)} \sum_{w \in WP_D} t^{\text{area}(D)} (-q)^{\ell(\alpha) - \ell(\beta)} q^{\text{dinv}(w) + \text{doff}_\beta(D) - \text{doff}_{\overleftarrow{\alpha}}(DP(\overleftarrow{\beta}))} x^w. \end{aligned}$$

For a Dyck path D in the sum, let $\gamma = \text{touch}(D)$. Note that $D \leq DP(\beta)$ implies that $\gamma \leq \beta \leq \alpha$, and thus we may rearrange sums as follows:

$$\begin{aligned} &\sum_{\beta \leq \alpha} \sum_{D \leq DP(\beta)} (-q)^{\ell(\alpha) - \ell(\beta)} q^{\text{doff}_\beta(D) - \text{doff}_{\overleftarrow{\alpha}}(DP(\overleftarrow{\beta}))} \\ &= \sum_{\gamma \leq \alpha} \sum_{\substack{D \\ \text{touch}(D)=\gamma}} \sum_{\substack{\beta \\ \gamma \leq \beta \leq \alpha}} (-1)^{\ell(\alpha) - \ell(\beta)} q^{\ell(\alpha) - \ell(\beta) + \text{doff}_\beta(D) - \text{doff}_{\overleftarrow{\alpha}}(DP(\overleftarrow{\beta}))}. \end{aligned}$$

Lemma 5.7 allows us to conclude that (5.14) reduces to Conjecture 4.5. \square

Corollary 5.10. *Conjecture 4.2 is true if and only if Conjecture 4.3 is true.*

REFERENCES

- [BDZ10] N. Bergeron, F. Descouens, and M. Zabrocki, *A filtration of (q, t) -Catalan numbers*, Adv. in Appl. Math. **44** (2010), no. 1, 16–36.
- [BGHT99] F. Bergeron, A. M. Garsia, M. Haiman, and G. Tesler, *Identities and positivity conjectures for some remarkable operators in the theory of symmetric functions*, Methods Appl. Anal. **6** (1999), 363–420.
- [GH01] A. M. Garsia and J. Haglund, *A positivity result in the theory of Macdonald polynomials*, Proc. Nat. Acad. Sci. U.S.A. **98** (2001), 4313–4316.
- [GH02] ———, *A proof of the q, t -Catalan positivity conjecture*, Discrete Math. **256** (2002), 677–717.
- [GXZ10] A. Garsia, G. Xin, and M. Zabrocki, *Hall-Littlewood operators in the theory of parking functions and diagonal harmonics*. International Mathematics Research Notices 2011; doi:10.1093/imrn/rnr060.
- [Hag03] J. Haglund, *Conjectured statistics for the q, t -Catalan numbers*, Adv. Math. **175** (2003), no. 2, 319–334.
- [Hag04] ———, *A proof of the q, t -Schröder conjecture*, Internat. Math. Res. Notices **11** (2004), 525–560.
- [Hag08] James Haglund, *The q, t -Catalan numbers and the space of diagonal harmonics*, University Lecture Series, vol. 41, American Mathematical Society, Providence, RI, 2008, With an appendix on the combinatorics of Macdonald polynomials.
- [Hai01] M. Haiman, *Hilbert schemes, polygraphs, and the Macdonald positivity conjecture*, J. Amer. Math. Soc. **14** (2001), 941–1006.

- [Hai02] ———, *Vanishing theorems and character formulas for the Hilbert scheme of points in the plane*, Invent. Math. **149** (2002), 371–407.
- [HHLRU05] J. Haglund, M. Haiman, N. Loehr, J. B. Remmel, and A. Ulyanov, *A combinatorial formula for the character of the diagonal coinvariants*, Duke J. Math. **126** (2005), 195–232.
- [Hic10] A. Hicks, *Two parking function bijections refining the q, t -Catalan and Schröder recursions*, preprint, 2010.
- [Jin91] Nai Huan Jing, *Vertex operators and Hall-Littlewood symmetric functions*, Adv. Math. **87** (1991), no. 2, 226–248.
- [Lam08] T. Lam, *Schubert polynomials for the affine grassmannian*, J. Amer. Math Soc **21** (2008), no. 1, 259–281.
- [LLM03] L. Lapointe, A. Lascoux, and J. Morse, *Tableau atoms and a new Macdonald positivity conjecture*, Duke Math. J. **116** (2003), no. 1, 103–146.
- [LM03] L. Lapointe and J. Morse, *Schur function analogs for a filtration of the symmetric function space*, J. Combin. Theory Ser. A **101** (2003), no. 2, 191–224.
- [LM07] L. Lapointe, , and J. Morse, *A k -tableaux characterization of k -schur functions*, Adv Math **213** (2007), no. 1, 183–204.
- [LM08] L. Lapointe and J. Morse, *Quantum cohomology and the k -Schur basis*, Trans. Amer. Math. Soc. **360** (2008), no. 4, 2021–2040.
- [LW08] N. Loehr and G. Warrington, *Nested quantum Dyck paths and $\nabla(s_\lambda)$* , Int. Math. Res. Not. IMRN **5** (2008), Art. ID rnm 157, 29.
- [Mac95] I. G. Macdonald, *Symmetric functions and Hall polynomials*, Oxford Mathematical Monographs, second ed., Oxford Science Publications, The Clarendon Press Oxford University Press, New York, 1995.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PENNSYLVANIA, PHILADELPHIA, PA 19104-6395

E-mail address: jhaglund@math.upenn.edu

DEPARTMENT OF MATHEMATICS, DREXEL UNIVERSITY, PHILADELPHIA, PA 19104

E-mail address: morsej@math.drexel.edu

DEPARTMENT OF MATHEMATICS AND STATISTICS, YORK UNIVERSITY, TORONTO, ONTARIO M3J 1P3, CA

E-mail address: zabrocki@mathstat.yorku.ca